

For complete warnings and directions, see blister card packaging.

Manufactured by Wisconsin Pharmacal Company, LLC, Jackson, WI 53037 USA

Distributed by Spectrum, Division of United Industries Corporation

PO Box 142642, St. Louis, MO 63114-0642

Questions or Comments? Call 1-800-767-9927. 17-15630 ©2014 UIC

BiteMD

External Analgesic
First Aid Antiseptic



from the
makers of

Cutter
BRAND

Net 0.5 fl oz (14.7 mL)

Active Ingredients: Benzocaine 5%, Camphor 3%, Phenol 1.35%

DIRECTIONS: Adults and children 2 years and over – apply to bite area by pressing stick straight down repeatedly until liquid flows, then dab sparingly. If itching persists, apply again in 15 minutes. Apply to affected area 1 to 3 times daily. For ticks and bees, remove tick or stinger before treatment.

WARNING: FOR EXTERNAL USE ONLY. • Do not get in eyes.

BiteMD[®]

External Analgesic
First Aid Antiseptic



from the makers of

Cutter
BRAND

INSECT BITE RELIEF

- ✓ **STOPS PAIN**
- ✓ **RELIEVES ITCHING**
- ✓ **REDUCES INFECTION**



Chiggers Wasps Mosquitoes Ticks



Soft Tip For
Controlled Application

Net 0.5 fl oz (14.7 mL)




BiteMD[®]

External Analgesic
First Aid Antiseptic



from the makers of
Cutter
BRAND



BiteMD, when applied soon after an insect bite, begins to work immediately to counteract the effects of bites and stings from bees, wasps, mosquitoes, ticks, hornets, ants, fire ants, deerflies, chiggers, sand fleas and horseflies.

DRUG FACTS

Active Ingredients:

Benzocaine 5% Analgesic, Anesthetic, Antipruritic
Camphor 3% Topical Analgesic
Phenol 1.35% First Aid Antiseptic

Purpose:

Uses: First aid for temporary relief of: ■ Pain ■ Itching ■ Infection

Warnings:

For external use only

■ Do not get in eyes.

When using

this product

■ Do not apply over large areas of the body or bandage.
■ In case of deep or puncture wounds, animal bites or serious burns, consult a doctor.

Stop use and ask doctor if condition worsens; or if symptoms persist for more than 7 days, clear up and occur again within a few days. Do not use longer than one week unless directed by doctor.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If in eyes, flush with water for 15 minutes and call a physician.

Directions

- Clean area.
- Apply to the bite area by pressing straight down repeatedly until liquid flows, then dab sparingly.
- If itching persists, apply again in 10 to 15 minutes.
- For ticks and bees, remove tick or stinger before treatment.

Other Information

■ Store at room temperature.

Adult and children 2 years and over

Apply to affected area 1 to 3 times daily

Children under 2 years

Consult a doctor

Inactive Ingredients

Propylene Glycol

Questions or Comments? Call 1-800-767-9927.

IMPORTANT: Read all information on this packaging and keep for future reference. This contains information not included on the stick label.

Manufactured by

Wisconsin Pharmacal Company, LLC

Jackson, WI 53037 USA

Distributed by Spectrum

Division of United Industries Corporation

PO Box 142642, St. Louis, MO 63114-0642

16-11359 ©2014 UIC

FPO UPC
0 71121 95614 9